

ProDiGyA - Interim Analysis of a Real-world, Prospective, Non-interventional Study Evaluating Daridorexant in Patients with Insomnia Disorder

Yaroslav Winter¹, Jens Kuhn², Göran Hajak³, Stefan Ries⁴, Corinna Frohn⁵, Cornelius Bachmann⁶, Eva Kugler⁷, Julia Kleylein-Sohn⁷, Michael Saletu⁸, Lucie Bartova⁹

1. Department of Neurology, UKS – Saarland University Hospital, Homburg, Germany | 2. Specialized Clinic for Psychiatry, Psychotherapy, and Neurology, Alexianer Köln GmbH & Department of Psychiatry and Psychotherapy, University Hospital Cologne, Germany | 3. Munich Psychiatry Service Professor Hajak at ProSomno Clinic, Munich, Germany | 4. Neurological-Psychiatric Group Practice, Neuro-Centrum Odenwald, Germany | 5. MVZ Solingen-Aufderhöhe, Diakonie Bethanien gGmbH, Solingen, Germany | 6. SomnoDiagnostics, Osnabrück, Germany | 7. Idorsia Pharmaceuticals Germany GmbH, Munich, Germany | 8. Sleep center at State Hospital Graz II Department of Neurology Austria, Sleep Society, Austria & Mannswoth Schwechat Medical Center, Schwechat-Mannswoth, Austria | 9. Department of Psychiatry and Psychotherapy (Clinical Division of General Psychiatry), Medical University of Vienna, Austria & Wiener Couch, Vienna, Austria

W130

Introduction

- **Chronic insomnia** affects ~6% of adults in Germany¹
- **Daridorexant** is a dual orexin receptor antagonist, effective on both sleep and daytime functioning as demonstrated in large clinical trials in adults with chronic insomnia²
- **Real-world data** for daridorexant remain **limited**
- **ProDiGyA** evaluates effectiveness and safety in routine practice

Study objective

To generate real-world evidence on the patient characteristics, treatment patterns, effectiveness and safety associated with daridorexant use in adults with chronic insomnia in Germany and Austria. The primary endpoint of this study is the maximum change in ISI score from baseline during the observation period of up to 12 months.

Results

Table 1. Chronic insomnia rarely comes alone and often remains untreated


Characteristic	
Female, n(%)	126 (55%)
Mean age (SD)	54.0 (15.6)
Insomnia duration ≥ 1 year, n(%)	190 (83%)
Most common insomnia symptom*: difficulty staying asleep, n(%)	203 (89%)
≥ 1 comorbidity, n (%)	193 (85%)
Prior insomnia pharmacotherapy, n (%)	113 (50%)

Patients: n=228; sites: n=28; *Multiple answers possible

85% had ≥ 1 comorbidity: 45% depression, 20% RLS and 20% OSA

83% had chronic insomnia ≥ 1 year: 51% 1-4 year, 20% 5-10 years, 13% > 10 years

Figure 2. Daridorexant is often used as first-line pharmacotherapy




50% were pharmacotherapy naïve

Dominant previous pharmacological therapy were sedating antidepressants


Only 3% had prior CBT-I

Figure 4. Insomnia severity improved early and continued to improve over time



Primary endpoint (interim analysis): Maximum mean change from baseline in ISI: -6.35 points (SD 5.66)

Figure 5. Reductions in depressive symptom scores were observed over time in patients with comorbid depression



Data are shown for the subgroup with comorbid depression only. Data are shown as mean with standard deviation



References

1. Schlack, R. et al. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz. 2013;56(5-6):740-8.
2. Mignot et al. *Lancet Neurol*. 2022 Feb;21(2):125-139
3. SmPC Daridorexant, current version

Disclosures


The study was funded by Idorsia Pharmaceuticals Germany GmbH. Conflict of interest (relevant for this poster): YW, JK, GH, SR, CF, MS, LB are investigators of the study and received speaker/consultant fees from Idorsia; EK, JKS are employees of Idorsia

Poster presented at the Sleep Europe Congress 2026

Methods

Interim analysis includes 228 patients with data available up to month 3

- Prospective, non-interventional, multicenter study in Germany and Austria
- Visit schedule:



- Daridorexant-naïve adults with chronic insomnia according to label³
- Interim analysis includes baseline (n=228), month 1 (n=145) and month 3 data (n=29).
- Data cut-off: 6 March 2026; only German sites included (28 sites).
- In total 800 patients to be enrolled, last-patient-out planned November 2027

Figure 1. Pre-treated patients were switched to Daridorexant mainly via stop & start





Table 2. No new safety issues were observed in this interim analysis

- Safety profile consistent with previous evidence
- No new safety issues were observed
- Low number of serious adverse events and AESIs were reported

Safety outcome	Patients, n(%)
Patients with ≥TEAE	38 (16.6)
Patients with ≥TEADR	26 (11.4)
Patients with ≥TESAE	2 (0.9)
Patients with ≥TEAESI	2 (0.9)

TAESI: Treatment-emergent adverse event of special interest; TE(S)AE: Treatment-emergent (serious) adverse event; TEADR: Treatment-emergent adverse drug reaction

Figure 3. Daridorexant was mainly used as 50 mg monotherapy with high adherence




78% started with 50 mg Daridorexant

80% received Daridorexant monotherapy

Daily intake was reported by 87% at V1

10% discontinued treatment

Conclusions



Patient population

Chronic insomnia rarely comes alone and often remains untreated



Treatment use

Daridorexant was mainly used as 50 mg monotherapy with high adherence



Effectiveness

Clinically meaningful improvements in insomnia severity were observed



Safety

No new safety issues were observed in this interim analysis

These interim data support the effectiveness of daridorexant in a broad real-world population treated in routine clinical practice with no new safety issues.